

In the format provided by the authors and unedited.

Shared strategies for β -lactam catabolism in the soil microbiome

Terence S. Crofts^{1,2}, Bin Wang^{1,2}, Aaron Spivak², Tara A. Gianoulis^{3,10}, Kevin J. Forsberg^{2,8}, Molly K. Gibson², Lauren A. Johnsky⁴, Stacey M. Broomall⁴, C. Nicole Rosenzweig⁴, Evan W. Skowronski^{4,9}, Henry S. Gibbons⁴, Morten O. A Sommer⁵ and Gautam Dantas^{1,2,6,7*}

¹Department of Pathology and Immunology, Washington University in St Louis School of Medicine, Saint Louis, MO, USA. ²The Edison Family Center for Genome Sciences and Systems Biology, Washington University in St Louis School of Medicine, Saint Louis, MO, USA. ³Wyss Institute for Biologically Inspired Engineering, Harvard, Cambridge, MA, USA. ⁴US Army Edgewood Chemical Biological Center, Aberdeen Proving Ground, Aberdeen, MD, USA. ⁵Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Lyngby, Denmark. ⁶Department of Molecular Microbiology, Washington University in St Louis School of Medicine, Saint Louis, MO, USA. ⁷Department of Biomedical Engineering, Washington University in St Louis, Saint Louis, MO, USA. ⁸Present address: Division of Basic Sciences, Fred Hutchinson Cancer Center, Seattle, WA, USA. ⁹Present address: TMG Biosciences, LLC, Austin, TX, USA. ¹⁰Deceased. *e-mail: dantas@wustl.edu

Supplementary table 1. Description of previously selected antibiotic-catabolizing soil isolates

Isolate name ^a	Prior designation	Organism	Initial selection	β-lactam catabolism	
				Penicillin	Carbenicillin
ABC02	CA-S3F-1	<i>Paraburkholderia caledonica</i> CA-S3F-1	Carbenicillin	+	+
ABC07	PE-S1G-1	<i>Pseudomonas</i> sp. PE-S1G-1	Penicillin	+	-
ABC08	PE-S2R-1	<i>Pandoraea</i> sp. PE-S2R-1	Penicillin	+	-
ABC10	PE-S2T-3	<i>Pandoraea</i> sp. PE-S2T-3	Penicillin	+	-

^aData, isolate selection and growth described in Dantas *et al* (2008)¹⁵

Supplementary table 2. Total β-lactamase and β-lactamase-like genes identified in ABC strain genomes

Ambler class	ABC02	ABC07	ABC08	ABC10	Total
Class A	1	0	0	0	1
Class B	0	0	0	0	0
Class C	5	2	4	4	15
Class D	0	0	1	1	2
Incomplete annotation*	3	8	3	3	17
Total	9	10	8	8	35

*Includes β-lactamases without Ambler class identification, and putative β-lactamases annotated as serine proteases or, predominantly, MBL fold metallo-hydrolases


Supplementary table 3. β-lactamases up-regulated by growth on penicillin in ABC strain genomes

Ambler class	ABC02	ABC07	ABC08	ABC10	Total
Class A	1	0	0	0	1
Class B	0	0	0	0	0
Class C	1	2	1	1	5
Class D	0	0	1	1	2
Total	2	2	2	2	8

Supplementary table 4. Primer sequences


Primer name	Sequence	Purpose
5093BW	GTCGAC TCTAGA ATGAGTATTGTTGAACTGGATGC	Amplification of <i>put1</i> gene for cloning into pET28B vector
5094BW	CCAGTGAA CTCGAG TCAACCGGTCGCCGAGAGGC	Amplification of <i>put1</i> gene for cloning into pET28B vector
5095BW	GTCGAC TCTAGA ATGAGCCCACACCAGCACATCC	Amplification of <i>put2</i> gene for cloning into pET28B vector
5096BW	CCAGTGAA CTCGAG TCAGGCGGGCTTGAGCCA	Amplification of <i>put2</i> gene for cloning into pET28B vector
54TSC	CGAGATCTCGATCCCGCG	Diagnostic PCR for cloning of inserts into pET28B vector
55TSC	CTAGTTATTGCTCAGCGGTGG	Diagnostic PCR for cloning of inserts into pET28B vector
5777TSC	ATGGGTACCATGAAAAATAGAAAT CGTATGATCGTG	Amplification of <i>E. coli</i> penicillin G amidase gene for cloning into pZA11 vector
5779TSC	ATGCTGCAGTTATCTCTGAACGTGC AACACTTC	Amplification of <i>E. coli</i> penicillin G amidase gene for cloning into pZA11 vector
5780TSC	TTCCTGCAGCCCGG	Amplification of pZA11 vector by inverse PCR
5781TSC	CCCGGTACCTTTCTCC	Amplification of pZA11 vector by inverse PCR
22TSC	GATACTGAGCACATCAGCAGGA	Diagnostic PCR for cloning of inserts into pZA11 vector
5723TSC	GGCATCTTCCAGGAAATCTCC	Diagnostic PCR for cloning of inserts into pZA11 vector
6057TSC	atgttacgcagcagcaacg	Diagnostic PCR for gentamicin resistance cassette
6058TSC	ttaggtggcggtacttggg	Diagnostic PCR for gentamicin resistance cassette
6059TSC	atgtccaccaacttatcagtataaag	Diagnostic PCR for <i>tetA</i>
6060TSC	tcagcgatcggtctg	Diagnostic PCR for <i>tetA</i>
6032TSC	gtcgactctagaggatccccGCTGTGGAGAGG CCGGCT	Amplification of ~1,000 bp upstream of <i>paaF</i>
6033TSC	agttggtggacatGAGTTCACTCGAATTAT TTTTGTGTATCAGCCCAC	Amplification of ~1,000 bp upstream of <i>paaF</i>
6034TSC	gagtgaactATGTCCACCAACTTATCA GTGATAAAG	Amplification of <i>tetA</i> for <i>paaF</i> allelic replacement
6035TSC	ggtgcagcagTCAGCGATCGGCTCGTTG	Amplification of <i>tetA</i> for <i>paaF</i> allelic replacement
6036TSC	cgategetgaCTGCTGCACCTTTTGGCC C	Amplification of ~1,000 bp downstream of <i>paaF</i>
6037TSC	ccgaattcgagctcgagcccTTGCGCCTGGGT	Amplification of ~1,000 bp

	GCCCTG	downstream of <i>paaF</i>
6087TSC	tgtaaagcaagcttctgcaggtcgactctagaggatcccc CCTCCAGCGCCCCCGGCC	Amplification of ~1,000 bp upstream of penicillin catabolon
6088TSC	actgataagttggtggacatGGTGATTCTCCTC GCAGGCGGGTG	Amplification of ~1,000 bp upstream of penicillin catabolon
6089TSC	cgctgcgaggagaatcaccATGTCCACCAAC TTATCAGTGATAAAG	Amplification of <i>tetA</i> for penicillin catabolon allelic replacement
6090TSC	agatacaggcccgtcgcggcTCAGCGATCGG CTCGTTG	Amplification of <i>tetA</i> for penicillin catabolon allelic replacement
6091TSC	ggcaacgagccgatcgctgaGCCGCGACGGG CCTGTAT	Amplification of ~1,000 bp downstream of penicillin catabolon
6092TSC	aattaaggtaccgaattcgagctcgagcccGACATA AAGACTCTTGGGGCAAAAAACCTG	Amplification of ~1,000 bp downstream of penicillin catabolon
6093TSC	GGGCTCGAGCTCGAATTCGG	Amplify pEXG2 plasmid for Gibson assembly to make ABC07 KOs
6031TSC	GGGGATCCTCTAGAGTCG	Amplify pEXG2 plasmid for Gibson assembly to make ABC07 KOs
6098TSC	ATGAACATGCCTATTCCAAACGCCG TG	Diagnostic PCR for <i>paaF</i>
6099TSC	TCATGCCTTAGGGCGACGATCC	Diagnostic PCR for <i>paaF</i>




Supplementary figure 1. ABC strains exhibit multi-class, cross-generational resistance to β -lactam antibiotics

ABC strains were grown in rich media with 1 g/l β -lactam antibiotics including 3rd and 4th generation penicillins (ampicillin/amoxicillin and piperacillin respectively), 2nd, 3rd, and 4th generation cephalosporins (cefuroxime, cefdinir/ceftriaxone/cefotaxime, and cefepime respectively), a monobactam (aztreonam) and two carbapenems (imipenem and meropenem). Average culture density measured by OD₆₀₀ of three cultures with standard deviation is shown.




Supplementary figure 2. ABC strains catabolize diverse substrates including aromatic compounds

Growth of four ABC strains were assayed on 190 different carbon sources. Only carbon sources for which growth exceeded OD₆₀₀ 0.05 for at least one strain are displayed. Aromatic carbon sources are boxed for emphasis.




Supplementary figure 3. Growth of ABC07, *paaF*, and *put* knock-outs on penicillin degradation products and other β -lactam antibiotics

Growth curves of wild-type ABC07 (black), and *paaF::tetA* (red) and *put::tetA* (blue) strains in minimal media containing (a) penicillin, (b) benzylpenicilloic acid, (c) phenylacetic acid, (d) 6-aminopenicillanic acid, (e) carbenicillin, (f) ampicillin, (g) cefuroxime, or (h) glucose as sole carbon source. Measurements are the average of triplicate cultures with standard error of the mean displayed.



Supplementary figure 4. Phylogenetic analysis of penicillin-responsive amidases, amidohydrolases, and hydrolases from ABC strains compared to penicillin amidases
Maximum Likelihood tree of penicillin amidase amino acid sequences (shown in blue), including *E. coli* penicillin amidase and penicillin-upregulated amidase, amidohydrolase, and hydrolase amino acid sequences from ABC strains (shown in red). Node support is given by bootstrap fraction.




Supplementary figure 5. *In vitro* characterization of ABC07 Put1 and Put2 from *E. coli* overexpression

(a) SDS-PAGE gel of his-tagged Put1 and Put2 following over expression in *E. coli* and purification by IMAC resin. Expected protein masses are approximately 54.4 kDa and 44.0 kDa for Put1 and Put2 respectively. (b through g) Enzyme kinetic analysis of Put1, Put2 (panels b and c only), commercial *E. coli* penicillin amidase, or no enzyme with (b) 6-nitro-3-(phenylacetamido)benzoic acid (NIPAB) (penicillin amidase K_M 70.12±16.13 μ M), (c) *p*-nitroacetanilide, (d) N-phenylacetyl-*p*-aminobenzoate (penicillin amidase K_M 68.97±24.45 μ M), (e) L-glutamate 1-(*p*-nitroaniline), (f) L-glutamate γ -(*p*-nitroaniline) (Put1 K_M 73.33±10.53 μ M), or (g) N α -benzoyl-DL-arginine *p*-nitroaniline (penicillin amidase K_M 1.99±1.93 mM) as substrates.


A

		*	#	
ABC07 Put1	132	QTYNSVFGTTGNAY-----	-----VMSLCESALKDFAAL	292
<i>E. coli</i>	280	ANGLAGYPTTSNMWVIGKSKA...	NPQSGYIANWNNSPQKDYPAS	540
<i>K. cryocrescens</i>	280	ANGLAGYPTTSNMWVIGKNKA...	NPQSGYIANWNNSPQKDYPAS	539
<i>B. megaterium</i>	256	EELGLPLKIGSNAAIVGSEKS...	NPKNGYVVWNNKPSKEWVN-	519
<i>A. viscosus</i>	256	EELGLPLKIGSNAAIVGSEKS...	NPKNGYVVWNNKPSKEWVN-	519
<i>Pseudomonas</i> sp.	230	GDASDAAGGGSNNWAVAPGRT...	DPPGGLIVTANNRVVADDHP-	490

B




C




Supplementary figure 6. Sequence and structural comparison of Put1 with penicillin amidase enzyme family

(a) Multiple sequence alignment of Put1 enzyme alongside penicillin G amidase homologs. Amino acid residues conserved among penicillin amidases and N-terminal nucleophile hydrolases, the catalytic serine (*) and the oxyanion hole (#), are highlighted. Overlays and fit (RMSD/aligned residues) of *E. coli* penicillin amidase with (b) the Phyre predicted Put1 three dimensional structure (6.79 Å/88 residues) and (b) penicillin G amidases homologs from *P. rettgeri* (1.71 Å/744 residues), and *K. citrophilia* (1.39 Å/552 residues).



Supplementary figure 7. Put1 hydrolyzes the amide bond of benzylpenicilloic acid but shows little degradative activity with penicillin

(a) LCMS analysis of penicillin degradation by *E. coli* penicillin amidase or Put1 by extracted ion count monitoring at 335 m/z corresponding to singly protonated penicillin. Within each condition traces represent (bottom to top) incubation for 30 min, 300 min, and 22.5 hr. (b) Put1 and *E. coli* penicillin amidase were incubated with penicillin or benzylpenicilloic acid as substrates overnight. Hydrolysis of an amide bond (the phenylacetamide bond in penicillin and benzylpenicilloic acid or the β -lactam ring in penicillin) was detected by derivatization of newly revealed amino groups by the reagent NBD-Cl followed by fluorescent quantification.



Supplementary figure 8. Test conditions for the expression and purification of Put1 from *E. coli*

SDS-PAGE gels of (a) insoluble and (b) soluble protein fractions from cell-free extracts of *E. coli* BL21(DE3) cells expressing *put1* on pET28B vector under the following conditions: TB brand MB induced with (1) no IPTG, (2) 100 μ M IPTG at inoculation, (3) 100 μ M IPTG at density, (4) 500 μ M IPTG at inoculation, (5) 500 μ M IPTG at density, and TB brand FB induced with (6) no IPTG, (7) 100 μ M IPTG at inoculation, (8) 100 μ M IPTG at density, (9) 500 μ M IPTG at inoculation, and (10) 500 μ M IPTG at density. For all conditions the majority of Put1 protein (MW 54.4 kDa) is found in the insoluble fraction.